

Initiating Aminoglycosides Safely

SPECIFIC AMINOGLYCOSIDES

- Gentamicin and Tobramycin (amikacin not discussed below, see [companion document](#))

SPECTRUM OF ACTIVITY

- Many gram-negative bacteria, including resistant Enterobacteriaceae and Pseudomonas spp.
- Synergistic activity with cell-wall agents against Gram-positive organisms, including staphylococci, streptococci and enterococci

INDICATIONS (Note: Use in combination with other antibiotics for empiric treatment. When possible, ALWAYS use an alternate agent to aminoglycosides for targeted treatment.)

- Febrile Neutropenia ([High-Risk](#) and [Solid Tumor/Lymphoma](#))
- Sepsis with potential for resistant pathogens
- Ventilator-associated pneumonia with septic shock
- True β -lactam allergy and treatment of:
 - [Intra-abdominal infections](#)
 - Gynecologic infections
 - [Complicated UTI/Pyelonephritis](#)
- [Surgical prophylaxis](#)
- Synergy for some Gram-positive infections (i.e. endocarditis due to organisms listed above)

CONTRAINDICATIONS

- Myasthenia Gravis
- Caution: known auditory or vestibular disease

DOSING: (Note: informed consent should be obtained for therapy of > 3 days duration)

1. Once daily dosing (ODD) preferred, contraindicated in:

- | | | |
|-----------------------|--------------------------------------|-------------------------|
| ○ Pregnancy | ○ Renal Failure
(eGFR < 40ml/min) | ○ Burns > 20% BSA |
| ○ Significant Ascites | ○ Surgical Prophylaxis | ○ Gram-positive Synergy |

Dose: 5mg/kg of actual body weight (ABW)*

Interval: generally q24h, give first dose and consult clinical pharmacist and see [companion document](#)

*Obesity: If >30% above ideal body weight (IBW), use Dosing Weight

IBW (Men): 50kg + 2.3kg(x inches above 60 in)
 IBW (Women): 45.5kg + 2.3kg(x inches above 60 in)

Dosing Weight = IBW + 0.4(ABW – IBW)

2. Multiple daily dosing (MDD)/conventional dosing – see [companion document](#)
3. Gram-positive synergy dosing (infective endocarditis) – see [companion document](#)

Disclaimer: This document is intended for internal use at Sinai Health System and University Health Network. Recommendations herein are based on existing literature and clinical practice and are subject to change at any time. Please refer to the [Terms and Conditions](#) for more details.