

Antibiotics should be withheld until sterile tissue cultures are obtained, unless the patient is septic or there is a concomitant soft tissue infection. Identification of the causative organism is very important in order to treat appropriately.

EVALUATION

- ✦ Obtain sterile tissue cultures by percutaneous aspirate or surgical deep culture (ideally before starting antibiotics)
- ✦ Wound swabs tend to reflect colonization, and often do not indicate the infecting organism at the site of osteomyelitis
- ✦ If septic, draw blood cultures

EMPIRIC CHOICE

- ✦ For diabetic foot infections, refer to the 1-page document on that topic
- ✦ Acutely Unwell or Septic:
 - ceftriaxone 2 g iv q24h +/- metronidazole 500 mg p.o./iv q12h (add metronidazole for sacral osteomyelitis) +/- vancomycin (if known to be colonized/previous infection with MRSA)
 - if known to be ESBL colonized/previous infection: Meropenem 1 g iv q8h +/- vancomycin if known to be colonized/previous infection with MRSA
- ✦ Not septic:
 - Await sterile culture results to guide treatment

DURATION

- ✦ 4-6 weeks
- ✦ Shorter courses could be considered if the infected bone has been appropriately debrided

ALTERNATIVES FOR ALLERGIES

- ✦ If septic: meropenem 1 g iv q8h (penicillin cross-reactivity is ~ 1%) +/- vancomycin
- ✦ Not septic: await sterile culture results to guide treatment

TOP ORGANISMS

- ✦ *Staphylococcus aureus*
- ✦ Streptococci
- ✦ Gram negative bacilli
- ✦ Anaerobes

IMMUNOCOMPROMISED HOST CONSIDERATION

- ✦ Same as for immunocompetent host

ADDITIONAL DIAGNOSTIC AND THERAPEUTIC COMMENTS

- ✦ X-ray be normal for the first 2 weeks of osteomyelitis; use CT or MRI for diagnosis if suspicion is high but x-ray is negative

Disclaimer: This document is intended for internal use at Sinai Health System and University Health Network. Recommendations herein are based on existing literature and clinical practice and are subject to change at any time. Please refer to the [Terms and Conditions](#) for more details.